

Provisional Report*⁾ on the

F.S. Heincke – cruise no HE 191

12.06. – 22.06. 2003

To the

Directorate of Fisheries, Norway (Fiskeridirektoratet)

Reference to

License for scientific research for natural resources on the Norwegian continental shelf etc. in the Norwegian part of the area applied for in the “Notifications of proposed research cruise” – Cruise HE 191

License no. 395/2003

received 07. April 2003 from Fiskeridirektoratet, Bergen, Norge

*) This report contains navigational data, detailed station lists, brief descriptions of used gear and sampling methods, and preliminary results on the analyses of the sampled material. Final results of the cruise cannot be presented in within six months after the cruise, but will be made available as soon as possible.

Report F.S. Heincke – cruise no HE 191

RV Heincke call sign: DBCK
Cruise number: HE 191
Dates of Cruise: 12.06.-22.06.2003
General subject of research: Geology/geochemistry, EU Project METROL (Methane fluxes in ocean margin sediments: microbiological and geochemical control)

Port Calls: Bremerhaven: 12.06.2003; 22.06.2003
Institute: Alfred Wegener Institute for Polar and Marine Research

Chief Scientist: Dr. Christian Borowski, MPI-Bremen, Celsiusstr. 1, D-28359 Bremen, Germany Tel. +49 421 2028 649, Fax +49 421 2028 580, e-mail cborowsk@mpi-bremen.de

1. Scientific crew, list and institute affiliation

1. Christian Borowski (chief scientist) MPI
2. Tomas Wilkop (technician) AWI
3. Regina Usbeck (scientist) AWI/Fielax
4. Nina Knab (scientist) MPI
5. Jens Kallmeyer (scientist) MPI
6. Maren Nickel (scientist) MPI
7. Fabian Jacobi (student) MPI
8. Barry Cragg (scientist) Univ. Bristol
9. Fiona Brock (scientist) Univ. Bristol
10. Dan Secrieru (scientist) GeoEcoMar
11. Christian Hübscher (scientist) Univ. Hamburg
12. Sofie Gradmann (student) Univ. Hamburg

AWI = Alfred Wegener Inst. for Polar and Marine Research, Columbusstr., 27568 Bremerhaven, Germany

MPI = Max Planck Institute for Marine Microbiology, Celsiusstr. 1, Bremen, Germany
Univ. Bristol = University of Bristol, Dep. Earth Sciences, Queens Road, Bristol BS8 1RJ, Great Britain.

GeoEcoMar = National Institute for Marine Geology and GeoEcology, 23-25 Dimitrie Onciul St, 70318 Bucharest, Romania

Univ. Hamburg = Geophysical Institute, University of Hamburg, Bundesstr. 55, Hamburg, Germany

2. Research program

This cruise and the scientific work is part of the collaborative project METROL sponsored by the EU (METROL: METHane fluxes in ocean margin sediments: microbiological and geochemical contROL) lead by the MPI-Bremen. Project partners

are Germany, Norway, Denmark, UK, The Netherlands, Romania, and Ukraine. In the framework of this project the question is addressed how the methane turnover is regulated in shallow gassy sediments of the European continental margin.

Objectives of this cruise were shallow hydro acoustic mapping of subsurface gas plumes and sediment sampling for (i) the quantification of the microbial turnover of methane in gassy sediments as well as the characterisation of the geochemical conditions for the anaerobic methane oxidation; (ii) the characterisation and identification of microorganisms involved in the methane oxidation in aerobic and anaerobic sediment layers. As part of these investigations, characteristic organic molecules are to be identified which can be used as biomarkers for the anaerobic methane oxidising microorganisms. At sites with active methane seepage, additional water samples should give information on the aerobic oxidation of methane in the water column.

Target working area was the Skagerrak Trench, stretching from shallow areas on the southern slope in the Danish EEZ to the deepest parts of the trench in the Norwegian EEZ. The purpose of the cruise was primarily to collect the samples which will be analyzed in the home laboratories during subsequent months.

3. Scientific equipment and sample treatment

3.1 Seismic acquisition equipment

The shallow seismic survey was carried out with two instruments, a water gun and a boomer:

- Water gun: S15.02 water gun (Seismic Systems/Sodera, USA) – a pneumatic seismic source using compressed air as air guns do. Compressed air stored in the firing chamber (0.16 l per "shot") is used to propel a water jet that creates vacuum cavities. These emit a strong bubble-free high-frequency acoustic pulse when they implode due to surrounding hydrostatic pressure. The pulse has a broad spectrum and a middle frequency of around 1.2 kHz. Penetration depth of the signal is maximum 200 m.
- Boomer: 230-1 Uniboom (EG&G, USA). The boomer is towed behind the ship at the water surface. Reflected sound frequencies of 0.6-2.5 kHz are recorded by the streamer hydrophones.

3.2 Echosounders

The navigation echosounder of RV Heincke was used for the determination of the water depth. A "fish finder" and an Atlas DESO25 sediment echosounder (33 kHz; Seatronics, USA) were used for the detection of pockmarks in water depths down to 550m.

3.3 Sediment sampling

- Gravity corer: A large gravity corer equipped with a top weight of approx. 700 kg recovered sediment cores of maximum 5.5 m length. The cores were sub-sampled for the analyses of various geochemical and microbiological parameters (see below).
- Rumor lot: A small gravity corer that samples the top 50-100 cm of sediment.
- Multiple corer: A multiple corer (type of Barnett et al. 1985) was equipped with five coring tubes of 9.5 cm inner diameter and 60 cm length.
- Van Veen grab: A 1/10-m² Van Veen Grab was used for quickly “probing” the sediment type at certain locations.

3.4 Water sampling

- CTD/rosette. The Rosette was equipped with 9 Niskin-bottles.

3.5 Treatment of sediment samples

The gravity cores were cut in 1-m sections immediately upon retrieval, and sub-sampling of the core sections was done by, A) pushing out the core at one end with a large piston and subsampling from the core end with small coring tubes or syringes, B) cutting windows in the side of the core liner with a vibrating saw blade and immediately inserting small coring tubes or syringes.

The sub-samples served for the analyses of the following parameters:

CH₄
 SO₄²⁻
 H₂S
 HCO₃²⁻
 Density/porosity
 Fe/Si
 Volatile fatty acids
 Microbial lipids
 Isotopic δ¹³C CH₄ & CO₂
 δ D₂O
 H¹⁴CO₃²⁻-methanogenesis
 Acetate ¹⁴C-methanogenesis
 Anaerobic oxidation of methane
 Sulfate reduction rate
 AODC bacterial counts
 Bacterial/archaeal 16S analyses
 Fluorescence In Situ Hybridization

Samples obtained with the Rumor Lot and the Multi corer were used to analyze pore water gradients of sulfate and determinations of sedimentation rate in the upper sediment layers, which are usually washed out in gravity cores.

4. Cruise Narrative:

12 June 03

12:00 MEST Leave port of Bremerhaven, heading to “Roter Sand“ off Bremerhaven in the German Bight.

13:06 MEST Start of scientific work with a CTD-rosette station at „Roter Sand“. Ship station 762

13:20 MEST Continue the cruise with heading to the Skagerrak. During the transit, installation of laboratories and scientific equipment (Gravity corer, Multi corer, Rumohr lot)

13-June-03

14:40 UTC Arrival in working area 1. Ship station 763: Test of Multi corer

15:15 UTC Test of seismic equipment at Pos. 57°47`N, 9°43`E

16:39 UTC Ship stations 764, 765: seismic profiles 1 and 2 with Water gun.

14-June-03

08:00 UTC End of seismic profile

11:06-17:10 UTC station work, ship stations 766-774: gravity corer and Rumohr lot, wind W 5.

17:10 UTC Seismic profile 3 with Water gun, ship station 775.

15-June-03

04:00 UTC End of profile, Pos. 58°19,3`N, 9°30,0`E

07:00 UTC Station work with Rumohr lot and Gravity corer, ship stations 776-781. Wind calms down to N 3

10:50-14:20 UTC profile with echosounder DESO25, ship station 782

15:06 UTC Station work at 58° 02,7`N, 9° 37,7`E, Gravity corer and CTD. Ship stations 783, 784

moving to starting point of seismic profile 4

20:40 UTC Seismic profile 4 with Water gun, ship station 785

16-June-03

02:19 UTC End of seismic profile 4, Pos. 58°05,3`N, 8°49,5`E

06:03 UTC Start of station work at pos. 58°03,3`N, 9°36,5`E alternating deployments of CTD/rosette, Multi corer, Rumohr lot, Gravity corer, and Van Veen grab. Ship stations 786-810.

Moving to starting point of seismic profile 5 at pos. 57°54,0`N, 9°46,9`E.

19:18 UTC Seismic profile 5 with Boomer, ship station 810-a

18-Jun-03

02:18 UTC End if profile 5, Pos. 58°08,5`N, 9°28,5`E

04:32 UTC Start with station work at Pos. 57°55,2`N, 9°45,3`E, alternating deployments of Rumohr lot, Gravity corer, CTD/rosette at various positions, ship stations 811-823

Moving to starting point of seismic profile 6 at pos. 58°03,5`N, 9°48`E

18:36 UTC Start with seismic profiles 6-16 with Water gun. Ship stations 824-834

18-June-03

06:28 UTC End of profile 16 at pos. 58°01,0`N, 9°31,8`E, moving to next position.

07:42 UTC Start with station work at pos. 57° 57,1`N, 9° 42,4`E.

Alternating deployments of Gravity corer, Rumohr lot, Van Veen grab, ship stations 835-847.




Fig 1. Cruise Track HE 191, with working areas 1 and 2.

15:55-16:15 UTC working-boat maneuver at 57° 57,1`N, 9° 42,4`E

16:45 UTC Transit to working area 2

18:54 UTC Pos. 58° 22,0`N, 10° 00,0`E

Start with seismic profiles 17-20 with water gun, ship stations 848-852

19-June-03

05:51 UTC End of Seismic profile 20, 58° 28,0`N, 9°43,9`E
moving to pos. 58° 27,7`N, 9°50,6`E

08:21 UTC Start with echosounder profiles with DESO25, ship stations 852-856

12:10 UTC Start with station work at pos. 58° 26,5`N, 9°49,0`E.

Alternating deployments of Gravity corer, Rumohr lot. Ship stations 857-861.




Fig 2.: Detailed cruise track in the Skagerrak Area.

- 16:25 UTC Wind picks up to WSW 7/8. End of station work.
 17:00 UTC Moving to position 58°16'N, 10°00'E.
- 20-June-03
 04:00 UTC On pos. 58°16'N, 10°00'E. Wind has not calmed (W 7/8), station work impossible
 07:00 UTC Secure and tie down scientific equipment
 08:45 UTC Pos. 57° 56.6'N, 9°43.4'E, Wind continuously speeds up. End of scientific work. Disassemble scientific equipment on deck. Return to Bremerhaven.
- 21-June-03
 Transit to Bremerhaven. Disassemble laboratory equipment.
- 22-June-03
 00:54 MEST Arrival in Bremerhaven

5. Scientific report and first results




Fig. 3: Seismic profiles in working areas 1 (#1-16) and 2 (#17-20).

5.1. Seismics

The high-resolution multi-channel seismic (MCS) data illustrate the correlation of fluid escape structures (e.g., pockmarks) on the seafloor with reflection pattern from both Quaternary and Mesozoic successions, which are typical of gas charged strata. Signal attenuation and acoustic whitening within the Quaternary succession of the upper slope indicate the presence of gas as already described by other authors (Bøe et al. 1998, Rise et al. 1999). The alignments of elongate depressions correlate with the location of bright reflections within Cretaceous strata which subcrop at the erosional unconformity beneath the Quaternary succession. The observations support the thesis that thermogenic gas from Cretaceous strata contributed to the seepage at the elongate depressions. MCS data reveal acoustic turbulence at and beneath the unconformity downslope of the edge of gas front in water depths of more than 400 m, which indicates the presence of gas. We suggest that gas accumulates beneath the unconformity beneath a Quaternary capping sequence. In the central Norwegian Trench pockmark lineaments lie above subcropping bright reflections within the Jurassic bedrock. Some pockmark lineaments lie above elongated near vertical faults in the overburden.




Fig 4: Overview on the sediment coring and CTD stations.

5.2. Sediments

The sediment sampling program primarily focused on cores along the depth gradient of the southern slope of the Skagerrak trench. Most of the cores were obtained along the seismic profile 1 (working area 1, see figs 1, 3 and 4, tabs 1 and 2) in water depths between 85 m and 540 m, while two gravity cores plus the accompanying Rumohr lot cores were sampled around 58°26.70' N and 9°50' E in approx 500 m water depth (working area 2). On board analyses concentrated on the determination of methane concentrations in the porewaters of the gravity cores, while all other biogeochemical and microbiological parameters will be analyzed during the subsequent months in the home laboratories. One major target on board was to identify the sediment depth of the sulfate methane transition zone (SMR) - the sediment layer in which the anaerobic oxidation of methane takes place, which is an important process for cycling of methane in marine sediments.

Area 1:

On the southern slope of the Skagerrak Trench, porewater methane was present in sediments between 85 m and 400 m water depth, while no methane occurred in the cores sampled at 540 m. In the shallower parts, methane was found below 2 m sediment depth, while at some deeper stations the SMR reached the top end of the cores. A transect through a pockmark showed an asymmetrical pattern of the methane distribution. Detailed analyses the SMR will reveal the associated geochemical and microbiological processes at the various sampling locations. Active methane seepage was explored at locations where such phenomena have been reported before (Dando et al. 1994, Zimmermann et al. 1999), but signals from profiles with the DESO15 echosounder could not be unequivocally interpreted and multi corer deployments remained unsuccessful. However, black gassy surface sediment was once recovered from a non-pockmark location with a Van Veen grab at 58°0.4' N and 9°31.64' E.

Area 2:

Pockmarks described by Rise et al. (1999) were the targets in area 2. Gravity core #857 was positioned in an area of suspected pockmark accumulation according to the multi channel seismics, while the apart positioned core #860 served as a control. Both cores did not contain dissolved methane, and it was decided not to further concentrate on this area.

Literature:

- Barnett PRO, Watsen J, Conelly D 1984. A multiple corer for taking virtually undisturbed samples from shelf, bathyal, and abyssal sediments. *Oceanologica Acta*, 399-408.
- Bøe R, Rise L, Ottesen D 1998. Elongate depressions on the southern slope of the Norwegian Trench (Skagerrak): morphology and evolution. *Marine Geology* 146, 191-203.
- Dando PR, Bussmann I, Niven SJ, O'Hara SCM, Schmaljohann R, Taylor L. J. 1994. A methane seep area in the Skagerrak, the habitat of the pogonophore *Siboglinum poseidoni* and the bivalve mollusc *Thyasira sarsi*. *Marine Ecology Progress Series* 107, 137-167.
- Rise L, Sættem J, Fanavoll S, Thorsnes T, Ottesen D, Bøe R 1999. Sea-bed pockmarks related to fluid migration from Mesozoic bedrock strata in the Skagerrak offshore Norway. *Marine and Petroleum Geology* 16, 619-631.
- Zimmermann S, Hughes RG, Flügel HJ 1997. The effect of methane seepage on the spatial distribution of oxygen and dissolved sulphide within a muddy sediment. *Marine Geology* 137, 149-157.

6. Station lists

Gear abbreviations:

- Seis multi channel seismics or boomer (only profile 5, station # 810)
- DESO DESO25 echosounder
- GC Gravity corer
- RL Rumor lot
- MUC Multi corer
- VV Van Veen grab
- CTD CTD/rosette

Tab 1: List of profiles

| Station # | PI | Gear | Profile # | Begin | UTC | Lat. (N) | Long. (E) | End | UTC | Lat. (N) | Long. (E) |
|-----------|----------|------|-----------|------------|----------|-------------|------------|------------|----------|-------------|------------|
| 764 | Hübscher | Seis | 1 | 13/06/2003 | 16:39:00 | 57°48.94764 | 9°54.22752 | 14/06/2003 | 0:15:00 | 58°16.0098 | 9°17.54076 |
| 765 | Hübscher | Seis | 2 | 14/06/2003 | 2:20:00 | 58°8.23944 | 9°31.55394 | 14/06/2003 | 7:43:48 | 57°50.41272 | 10°3.30498 |
| 775 | Hübscher | Seis | 3 | 00/01/1900 | 0:00:00 | ° | ° | 15/06/2003 | 3:44:00 | 58°28.0893 | 9°29.86722 |
| 782 | Usbeck | DESO | - | 15/06/2003 | 10:50:00 | 58°4.01076 | 9°41.50974 | 15/06/2003 | 14:20:20 | 58°3.00702 | 9°38.92692 |
| 785 | Hübscher | Seis | 4 | 15/06/2003 | 20:42:00 | 57°42.43182 | 9°0.191902 | 04/01/1900 | 2:19:00 | 58°5.01246 | 8°48.97194 |
| 810-a | Hübscher | Seis | 5 | 16/06/2003 | 19:18:00 | 57°0.968322 | 9°42.03702 | 17/06/2003 | 2:18:00 | 58°7.02462 | 9°33.18606 |
| 824-834 | Hübscher | Seis | 6-16 | 17/06/2003 | 18:36:00 | 58°3.1305 | 9°46.86636 | 18/06/2003 | 6:28:00 | 58°0.39828 | 9°32.5626 |
| 848 | Borowski | VV | - | 18/06/2003 | 13:37:25 | 58°4.04207 | 9°41.63519 | 18/06/2003 | 15:08:20 | 58°4.12856 | 9°41.63505 |
| 849 | Hübscher | Seis | 17 | 18/06/2003 | 18:54:00 | 58°22.1865 | 9°59.6532 | 18/06/2003 | 21:20:00 | 58°31.00842 | 9°43.87656 |
| 850 | Hübscher | Seis | 18 | 18/06/2003 | 22:12:00 | 58°30.98808 | 9°50.01708 | 18/06/2003 | 23:50:00 | 58°25.00902 | 10°0.00582 |
| 851 | Hübscher | Seis | 19 | 19/06/2003 | 1:07:00 | 58°30.00756 | 9°0.02898 | 19/06/2003 | 3:15:00 | 58°22.48596 | 9°48.45492 |
| 852 | Hübscher | Seis | 20 | 19/06/2003 | 4:09:00 | 58°21.98964 | 9°54.0219 | 19/06/2003 | 5:51:00 | 58°28.35912 | 9°0.727512 |
| 853 | Borowski | DESO | - | 19/06/2003 | 8:21:40 | 58°28.10497 | 9°50.83552 | 00/01/1900 | 0:00:00 | ° | ° |
| 854 | Borowski | DESO | - | 19/06/2003 | 0:00:00 | ° | ° | 00/01/1900 | 0:00:00 | ° | ° |
| 855 | Borowski | DESO | - | 19/06/2003 | 0:00:00 | ° | ° | 00/01/1900 | 0:00:00 | ° | ° |
| 856 | Borowski | DESO | - | 19/06/2003 | 0:00:00 | ° | ° | 19/06/2003 | 10:53:00 | 58°28.14376 | 9°50.59796 |

Tab. 2. Sediment stations.

| Date | Station # | PI | Gear | Gear on Bottom | | | Depth (m) |
|------------|-----------|----------|------|----------------|-------------|------------|-----------|
| | | | | UTC | Lat. (N) | Long. (E) | |
| 12/06/2003 | 762 | Usbeck | CTD | 13:06:30 | 53°50.69869 | 8°5.42066 | 13.3 |
| 13/06/2003 | 763 | Borowski | MUC | 14:42:00 | 57°46.36137 | 9°40.11117 | 0.0 |
| 14/06/2003 | 766 | Borowski | RL | 11:06:00 | 57°55.22302 | 9°45.319 | 85.9 |
| 14/06/2003 | 767 | Borowski | GC | 11:31:00 | 57°55.23984 | 9°45.31077 | 87.2 |
| 14/06/2003 | 768 | Borowski | RL | 12:51:01 | 58°1.08049 | 9°38.68776 | 265.2 |
| 14/06/2003 | 769 | Borowski | GC | 13:14:45 | 58°1.09333 | 9°38.682 | 266.6 |
| 14/06/2003 | 770 | Borowski | GC | 14:31:53 | 58°1.09535 | 9°38.62538 | 265.8 |
| 14/06/2003 | 771 | Borowski | RL | 15:19:20 | 58°1.47436 | 9°38.41335 | 269.4 |
| 14/06/2003 | 772 | Borowski | GC | 15:42:50 | 58°1.51573 | 9°38.45988 | 269.0 |
| 14/06/2003 | 773 | Borowski | RL | 16:38:54 | 58°2.2054 | 9°37.89611 | 309.0 |
| 14/06/2003 | 774 | Borowski | GC | 17:04:03 | 58°2.25072 | 9°37.97088 | 311.0 |
| 15/06/2003 | 776 | Borowski | RL | 6:51:58 | 58°2.80978 | 9°37.06143 | 354.9 |
| 15/06/2003 | 777 | Borowski | GC | 7:15:37 | 58°2.74319 | 9°36.85114 | 354.2 |
| 15/06/2003 | 778 | Borowski | RL | 7:59:29 | 58°3.23319 | 9°35.9924 | 391.7 |
| 15/06/2003 | 779 | Borowski | GC | 8:18:00 | 58°3.2183 | 9°36.03015 | 389.6 |
| 15/06/2003 | 780 | Borowski | RL | 9:21:40 | 58°6.4997 | 9°31.12237 | 537.4 |
| 15/06/2003 | 781 | Borowski | GC | 9:46:30 | 58°6.4911 | 9°31.08305 | 534.5 |
| 15/06/2003 | 783 | Borowski | GC | 15:06:50 | 58°2.70912 | 9°37.6833 | 339.2 |
| 15/06/2003 | 783 | Usbeck | CTD | 16:49:00 | 58°3.82266 | 9°41.87286 | 335.1 |
| 16/06/2003 | 786 | Borowski | GC | 6:03:35 | 58°3.3159 | 9°36.48907 | 386.4 |
| 16/06/2003 | 787 | Borowski | RL | 6:35:14 | 58°3.20124 | 9°36.2886 | 384.3 |
| 16/06/2003 | 788 | Borowski | RL | 7:01:00 | 58°3.31896 | 9°35.96484 | 393.4 |
| 16/06/2003 | 789 | Borowski | GC | 7:31:00 | 58°3.25332 | 9°36.00546 | 391.1 |
| 16/06/2003 | 790 | Borowski | GC | 8:31:00 | 58°3.93756 | 9°38.01942 | 394.4 |
| 16/06/2003 | 791 | Usbeck | CTD | 9:21:00 | 58°3.82776 | 9°41.86404 | 335.1 |
| 16/06/2003 | 792 | Usbeck | CTD | 9:48:40 | 58°3.82404 | 9°41.81622 | 335.1 |
| 16/06/2003 | 793 | Borowski | MUC | 10:14:00 | 58°3.81826 | 9°41.82409 | 335.5 |
| 16/06/2003 | 794 | Borowski | MUC | 10:56:00 | 58°3.8373 | 9°41.81444 | 335.4 |
| 16/06/2003 | 795 | Usbeck | CTD | 11:26:00 | 58°3.42601 | 9°41.75476 | 331.9 |
| 16/06/2003 | 796 | Borowski | VV | ??? | ° | ° | 334.8 |
| 16/06/2003 | 797 | Borowski | VV | 13:09:00 | 58°3.83406 | 9°41.86957 | 335.2 |
| 16/06/2003 | 798 | Borowski | VV | 13:25:04 | 58°3.86769 | 9°41.81056 | 337.4 |
| 16/06/2003 | 799 | Borowski | VV | 13:38:27 | 58°3.93611 | 9°41.71885 | 342.5 |
| 16/06/2003 | 800 | Borowski | VV | ??? | ° | ° | 0.0 |
| 16/06/2003 | 801 | Borowski | VV | 13:50:19 | 58°4.00206 | 9°41.64537 | 348.6 |
| 16/06/2003 | 802 | Borowski | VV | 14:01:30 | 58°4.06065 | 9°41.59228 | 350.9 |
| 16/06/2003 | 803 | Borowski | RL | 15:15:08 | 57°55.24644 | 9°45.2558 | 86.3 |
| 16/06/2003 | 804 | Borowski | RL | 15:21:20 | 57°55.2488 | 9°45.29531 | 86.2 |
| 16/06/2003 | 805 | Borowski | RL | 15:30:50 | 57°55.2731 | 9°45.28932 | 86.4 |
| 16/06/2003 | 806 | Borowski | RL | 15:40:00 | 57°55.25189 | 9°45.33377 | 86.2 |
| 16/06/2003 | 807 | Borowski | GC | 16:12:53 | 57°55.24532 | 9°45.33555 | 86.3 |
| 16/06/2003 | 808 | Borowski | GC | 16:51:17 | 57°55.27971 | 9°45.29464 | 86.3 |
| 16/06/2003 | 809 | Borowski | RL | 17:10:52 | 57°55.28502 | 9°45.34019 | 86.1 |
| 16/06/2003 | 810 | Borowski | RL | 17:23:50 | 57°55.26749 | 9°45.38729 | 86.9 |
| 17/06/2003 | 811 | Borowski | GC | 4:23:30 | 57°55.25827 | 9°45.29846 | 86.2 |
| 17/06/2003 | 812 | Borowski | GC | 4:51:19 | 57°55.24457 | 9°45.29993 | 86.1 |
| 17/06/2003 | 813 | Borowski | GC | 5:13:20 | 57°55.26239 | 9°45.31632 | 86.4 |
| 17/06/2003 | 814 | Borowski | GC | 7:04:50 | 57°55.23757 | 9°45.35759 | 85.7 |
| 17/06/2003 | 815 | Borowski | GC | 7:39:45 | 57°54.95186 | 9°45.17638 | 82.7 |
| 17/06/2003 | 816 | Borowski | GC | 10:35:15 | 57°57.12172 | 9°42.43051 | 146.8 |
| 17/06/2003 | 817 | Borowski | RL | 10:59:00 | 57°57.11617 | 9°42.42274 | 148.4 |
| 17/06/2003 | 818 | Borowski | RL | 11:09:30 | 57°57.11809 | 9°42.42783 | 148.1 |
| 17/06/2003 | 819 | Usbeck | CTD | 11:44:50 | 57°55.87234 | 9°44.94803 | 104.2 |
| 17/06/2003 | 820 | Borowski | GC | 16:01:00 | 58°2.22593 | 9°37.99144 | 307.7 |
| 17/06/2003 | 821 | Borowski | GC | 16:54:06 | 58°2.24558 | 9°37.98319 | 308.7 |
| 17/06/2003 | 822 | Borowski | RL | 17:15:20 | 58°2.25507 | 9°37.96951 | 309.0 |
| 17/06/2003 | 823 | Borowski | RL | 17:30:12 | 58°2.23817 | 9°37.96622 | 308.4 |
| 18/06/2003 | 835 | Borowski | GC | 7:42:08 | 57°57.12268 | 9°42.4473 | 149.2 |
| 18/06/2003 | 836 | Borowski | GC | 8:42:33 | 57°57.12532 | 9°42.45235 | 148.9 |
| 18/06/2003 | 837 | Borowski | RL | 9:03:28 | 57°57.12415 | 9°42.41944 | 147.9 |
| 18/06/2003 | 838 | Borowski | RL | 9:12:10 | 57°57.12071 | 9°42.44422 | 149.2 |
| 18/06/2003 | 839 | Borowski | GC | 10:15:00 | 57°56.91281 | 9°42.70767 | 120.2 |
| 18/06/2003 | 840 | Borowski | RL | 10:34:31 | 57°56.91388 | 9°42.68695 | 120.6 |
| 18/06/2003 | 841 | Borowski | RL | 10:46:56 | 57°56.92716 | 9°42.69726 | 121.0 |
| 18/06/2003 | 842 | Borowski | GC | 11:05:08 | 57°57.03626 | 9°42.56353 | 138.2 |
| 18/06/2003 | 843 | Borowski | RL | 11:18:12 | 57°57.02876 | 9°42.55548 | 137.9 |
| 18/06/2003 | 844 | Borowski | GC | 11:42:08 | 57°57.14379 | 9°42.39673 | 134.1 |
| 18/06/2003 | 845 | Borowski | RL | 11:53:30 | 57°57.15448 | 9°42.40029 | 132.0 |
| 18/06/2003 | 846 | Borowski | GC | 12:14:00 | 57°57.24911 | 9°42.2689 | 128.0 |
| 18/06/2003 | 847 | Borowski | RL | 12:23:20 | 57°57.24787 | 9°42.26087 | 128.0 |
| 19/06/2003 | 857 | Borowski | GC | 12:10:30 | 58°26.67075 | 9°49.65427 | 496.0 |
| 19/06/2003 | 858 | Borowski | RL | 14:25:07 | 58°26.69154 | 9°49.698 | 506.0 |
| 19/06/2003 | 859 | Borowski | RL | 14:49:24 | 58°26.68697 | 9°49.67996 | 504.0 |
| 19/06/2003 | 860 | Borowski | GC | 15:21:55 | 58°26.79149 | 9°50.03202 | 499.0 |
| 19/06/2003 | 861 | Borowski | RL | 16:25:00 | 58°26.76712 | 9°49.97556 | 493.0 |